

Infection Prevention and Control

Best Practice Guideline

Aerosol Generating Medical Procedures (AGMP)

A printed version of this guideline may not be the most recent version. The OFFICIAL version is located at <u>www.ipac.vch.ca</u>.

Site Applicability

Applies to all VCH healthcare settings.

Scope

All healthcare workers (HCW).

Purpose

To prevent transmission of infection associated with aerosols produced by aerosol generating medical procedures.

Background

Aerosol generating medical procedures (AGMP) are any procedure carried out on a patient/resident/client (referred to as "patient" in the remainder of the document) that can induce the production of aerosols of various sizes, including droplet nuclei. This is applicable to patients of all ages and in any care setting.

Medical procedures that generate aerosols or droplet nuclei in high concentration present a risk for opportunistic airborne transmission of pathogens not otherwise spread by the airborne route (e.g., SARS-CoV-2, Influenza) and increase the risk for transmission of organisms known to spread by the airborne route (e.g., TB).

The <u>BC AGMP Expert Working Group</u> assesses medical procedures to determine which ones are considered to be AGMPs based on review of current information and evidence. As per the expert working group, procedures are assessed as probable AGMP, possible AGMP, or non-AGMP reflecting the varying amount and quality of evidence currently available for each procedure. As new evidence emerges, the list will be reviewed and updated accordingly. For the purposes of this document and related VCH practice recommendations, procedures are classified as either AGMP or non-AGMP, see Table 1.

Point of care risk assessment

All healthcare workers (HCW) should perform a <u>point of care risk assessment</u> (PCRA) prior to an AGMP to select the appropriate personal protective equipment (PPE) and environmental controls.



Patients should be assessed for suspect or confirmed viral respiratory illness (VRI) or respiratory infection prior to performing an AGMP.

In an emergency situation where a complete assessment is not possible, if there is concern for infection consider using the highest level of protection (N95 respirator).

For AGMP in healthcare settings

- Limit the number of HCW in the room or patient care area (privacy curtains) to only those necessary for the procedure.
- HCWs should perform hand hygiene before donning and after removing PPE and on leaving the room/area.
- Eye and face protection (if worn) should be removed **after** leaving the room/area and disposed of in either a hands-free waste receptacle (if disposable) or in a separate receptacle to go for reprocessing (if reusable).
- In long-term care a private room for residents who require the use of AGMPs (e.g. CPAP/BiPAP/Nebulizers) is preferred, however, if a private room is not available place or maintain the resident in a multi-bed room and create physical barriers by drawing privacy curtains and/or closing doors when an AGMP is occurring.

Procedure

Clinical care situations that Require N95 Respirators ONLY When Performing AGMPs:

- Treating patients with known or suspected VRI, novel respiratory pathogen, or for whom status of respiratory infection is unknown (including novel/pandemic influenza, seasonal influenza, SARS-CoV-2).
- Performing irrigation on a wound with confirmed extrapulmonary tuberculosis.

Table 1: Aerosol Generating Medical Procedure Classification

Assessment and Diagnostic Procedures				
AGMP	Non-AGMP			
Autopsy	Arterial Blood Gas			
Bronchoscopy and Bronchoalveolar Lavage (BAL)	Cardiopulmonary exercise testing			
Cough reflex testing	Clinical (bedside) swallow screen/assessment			
Flexible/fiber optic endoscopic evaluation of voice and swallowing (FEES)	Videofluroscopic swallow study			
Methacholine challenge (i.e., bronchoprovocation test)	Overnight oximetry			
Nasopharyngeal aspirates	Nasopharyngeal scoping			
Nasopharyngeal washes	Spirometry			
Sputum induction				



Upper gastrointestinal endoscopy				
Therapeutic and Resuscitation Procedures				
AGMP	Non-AGMP			
CPAP and BIPAP	Airway suctioning (deep suction and open tracheal suctioning)			
CPR with bag mask ventilation	Chest compressions alone			
Intubation and extubation procedures*	Intranasal naloxone			
Direct Laryngoscopy	Transesophageal Echocardiogram (TEE)			
Non-heated nebulizer style of high flow oxygen	Heated High Flow Oxygen			
(single or double flow)**	(e.g., AIRVO, Optiflow)**			
Nebulized Therapy	Vibrating mesh nebulizer (e.g., Aerogen)			
Mastoidectomy	Low flow oxygen devices (e.g., nasal prongs, O ₂ mask, non-rebreather mask)			
Tracheotomy	Tracheostomy Care			

*Includes: Hyperoxygenation using nasal prongs before intubation, manual ventilation before intubation, breaking/opening closed ventilator circuits (intentionally or un-intentionally).

**Refer to Appendix A High Flow Oxygen Devices

AGMP Environmental Controls

- Whenever possible, AGMP should be performed in a private or procedure room with the door closed.
- When an N95 respirator is indicated, priority placement for a private or procedure room must be assessed prior to non-emergent AGMPs
 - Private room priority should consider: infectious respiratory status, frequency of AGMP indicated, and patient immune status. To establish private room priority, refer to <u>Bed</u> <u>Placement for Viral Respiratory Illness (VRI)</u>
 - When a private or procedure room is not available and the priority placement assessment has determined the AGMP will occur in place, draw the privacy curtains and remove any shared equipment, supplies or linens from the immediate vicinity prior to performing an AGMP.
- If the priority placement assessment is selected for an Airborne Infection Isolation Room (AIIR) or a private/procedure room, the room should remain vacant or an N95 respirator should continue to be worn until the <u>air settle/clearance time</u> has lapsed.

Changes to this document are subject to learnings, evidence, evaluation, and alignment with the <u>BC Provincial AGMP list</u>.



Infection Prevention and Control

References

- 1. Canadian Agency for Drugs and Technologies in Health (CADTH). (2011). Rapid Response Report: Systematic Review Aerosol-Generating Procedures and Risk of Transmission of Acute Respiratory Infections: A Systematic Review. Retrieved from: https://www.cadth.ca/media/pdf/M0023 Aerosol Generating Procedures e.pdf
- Center for Disease Control and Prevention. (2016). Prevention Strategies for Seasonal Influenza in Healthcare Settings: Guidelines and recommendations. Retrieved from: https://www.cdc.gov/flu/professionals/infectioncontrol/healthcaresettings.htm
- 3. OSHA, CDC, NIOSH. (2015). Hospital Respiratory Protection Program Toolkit: Resources for Respiratory Program Administrators. Retrieved from: <u>https://www.osha.gov/Publications/OSHA3767.pdf</u>
- Provincial Infection Control Network. (2020). 2019 Novel Coronavirus: Aerosol Generating Medical Procedures in Healthcare Settings. Retrieved from: <u>http://www.bccdc.ca/Health-Professionals-Site/Documents/2019nCoV_AGMP_PICNet.pdf</u>
- Provincial Infectious Diseases Advisory Committee (PIDAC). (2012). Routine Practices and Additional precautions in all healthcare Settings, 3rd edition. Retrieved from: <u>https://www.publichealthontario.ca/en/eRepository/RPAP_All_HealthCare_Settings_Eng2012.pdf</u>
- Public Health Agency of Canada. (PHAC). (2013). Routine Practices and Additional Precautions for Preventing the Transmission of Infection in Health care Settings. Retrieved from: http://publications.gc.ca/collections/collection 2013/aspc-phac/HP40-83-2013-eng.pdf
- 7. VCH. (2019). Respiratory Protection Standard. Retrieved from: <u>https://my.vch.ca/working-here-site/Documents/Respiratory-Protection-Standard.pdf</u>
- 8. World Health Organization. (2009). Natural Ventilation for Infection Control in Health Care Settings. Retrieved from: <u>http://apps.who.int/iris/bitstream/10665/44167/1/9789241547857_eng.pdf</u>
- 9. World Health Organization. (2007). Infection prevention and control of epidemic- and pandemic-prone acute respiratory infections in health care. Retrieved from: <u>http://apps.who.int/iris/bitstream/10665/112656/1/9789241507134_eng.pdf?ua=1</u>
- BC AGMP Expert Working Group. (2022). Aerosol Generating Medical Procedures: Medical procedures and Aerosol Generating Medical Procedures (AGMPs) used for patients with confirmed or suspected COVID-19. Retrieved from: <u>http://www.bccdc.ca/health-professionals/clinical-resources/covid-19-care/infectioncontrol/aerosol-generating-medical-procedures</u>



Appendix A: High Flow Oxygen Devices

	Non-Heated High Flow Oxygen Therapy	Heated High Flow Oxygen Therapy		
AGMP Classification	AGMP	Non-AGMP		
Product Used	Large Volume Nebulizers (Single or Double Flow)	AIRVO, Optiflow		
Product Differences	 Acts like a Ventolin nebulizer, except the liquid reservoir is much larger Nebulizer reservoir could become contaminated with patient secretions. No power cable Bubbling/Misting/Nebulization Oxygen supply comes directly from the wall or tank to the attached water reservoir, then to client 	 Air and oxygen move from the medical gas source to a blender, then over a heated container of sterile water which increases the relative humidity in the circuit Requires plugged in power cable No Bubbling/Misting/Nebulization Oxygen supply connected to heating device, then to client 		



Infection Prevention and Control

lung 15 2017					
June 15, 2017					
March 4, 2025					
March 4, 2025					
Allyson Hankins, Munira Murji, Jenna Horner, Edwin Tunod, Robyn Hunter,					
Michelle Danzer					
Allyson Hankins Date: March 4, 2025					
Description of Changes:		Revised By:	Effective		
			Date:		
Fixed broken link to BC AGMP Expert Working Group (pg. 1); Removed recommendation to open windows to improve ventilation (pg. 2); Removed section re: clinical situations that always require N95 respirator (pg. 2)		Allyson Hankins, Munira Murji, Jenna Horner, Edwin Tunod, Robyn Hunter, Michelle Danzer	March 4, 2025		
Added under - Clinical care situations that Require Respirators ONLY When Performing AGMPs Performing irrigation on a would with confirmed extrapulmonary tuberculosis (pg. 2)	N95	Allyson Hankins, Munira Murji, Jenna Horner, Edwin Tunod, Michelle Danzer	June 6, 2025		
	March 4, 2025 Allyson Hankins, Munira Murji, Jenna H Michelle Danzer Allyson Hankins Description of Changes: Fixed broken link to BC AGMP Expert Working Grou 1); Removed recommendation to open windows to improve ventilation (pg. 2); Removed section re: clin situations that always require N95 respirator (pg. 2) Added under - Clinical care situations that Require Respirators ONLY When Performing AGMPs Performing irrigation on a would with confirmed	March 4, 2025 Allyson Hankins, Munira Murji, Jenna Horner, Michelle Danzer Allyson Hankins Date: Description of Changes: Fixed broken link to BC AGMP Expert Working Group (pg. 1); Removed recommendation to open windows to improve ventilation (pg. 2); Removed section re: clinical situations that always require N95 respirator (pg. 2) Added under - Clinical care situations that Require N95 Respirators ONLY When Performing AGMPs Performing irrigation on a would with confirmed	March 4, 2025 Allyson Hankins, Munira Murji, Jenna Horner, Edwin Tunod, F Michelle Danzer Allyson Hankins Date: March 4, 2025 Description of Changes: Revised By: Fixed broken link to BC AGMP Expert Working Group (pg. 1); Removed recommendation to open windows to improve ventilation (pg. 2); Removed section re: clinical situations that always require N95 respirator (pg. 2) Allyson Hankins, Munira Murji, Jenna Horner, Edwin Tunod, Robyn Hunter, Michelle Danzer Added under - Clinical care situations that Require N95 Respirators ONLY When Performing AGMPs Performing irrigation on a would with confirmed extrapulmonary tuberculosis (pg. 2) Allyson Hankins, Munira Murji, Jenna Horner, Edwin Tunod,		