


VCH Infectious Disease Risk Screening Documentation in Cerner

Risk Factors for Antibiotic-Resistant Organisms

NOTE: VCH does not screen or isolate for VRE in any healthcare setting



If risk factor(s) identified, order and collect ARO swabs within 24 hours of admission:

Test to order	Risk Factor
MRSA & CPO	<ul style="list-style-type: none"> Healthcare outside of Canada within the last year Cerner will auto-order Contact Precautions
CPO	<ul style="list-style-type: none"> Household contact with known CPO in the last year Cerner will auto-order Contact Precautions
MRSA	<ul style="list-style-type: none"> All other risk factors listed

VCH Screening Swab Sites

*If MRSA & CPO screening indicated, collect one swab from each site(s) to test for both AROs

MRSA	CPO
<ul style="list-style-type: none"> Nares Perirectal or rectal/ostomy* If applicable: <ul style="list-style-type: none"> Open wound(s)* 	<ul style="list-style-type: none"> Rectal/ostomy (fecal stained)* If applicable: <ul style="list-style-type: none"> Sputum/tracheostomy if coughing Urine if catheterized Open wound(s)*

Travel History

Does not automatically initiate isolation or require screening swabs. Used for information purposes and in the event of an emerging disease.

Risk Factors and Symptoms/ARO Surveillance

Risk Factors and Symptoms/ARO Surveillance			
	Yes	No	Unable to obtain
*Fever			
*Diarrhea			
*Headache			
*Photophobia			
*Illness With Generalized Rash			
*New or Worsening Cough			
*Recent Exposure to Communicable Disease			
*History of AROs			
*History of CPO			
*Immunocompromised			

Unable to Obtain Current Visit Information

None

Clinical condition

Cognitive impairment

Language barrier

Physical impairment

Communicable Disease Exposed To:

Measles Chickenpox or shingles

Mumps Other:

If new or worsening cough, refer to site or organizational procedures.

Tuberculosis Status Description

Active pulmonary TB

Positive TB skin test

Treated

Recent exposure

Other:

Was the patient received as a transfer from another facility?

Not transferred

Transferred from ambulatory surgery center

Transferred from ED or observation unit of another hospital

Transferred from inpatient department or care facilities

Transferred from outpatient department or care facilities

Cerner will automatically order isolation based on risk factor(s) selected. If you select any of the following, [contact VCH Infection Prevention and Control \(IPAC\)](#)

Risk Factor	Automatic Isolation Order
History of AROs (only select for MRSA)	Contact Precautions <ul style="list-style-type: none"> Contact IPAC if patient reports an ARO history not previously known/flagged in Cerner
History of CPO	Contact Precautions <ul style="list-style-type: none"> Contact IPAC if patient reports a CPO history not previously known/flagged in Cerner
Diarrhea	Contact Plus Precautions
Recent Exposure to Communicable Disease, Measles	Airborne Precautions
Recent Exposure to Communicable Disease, Mumps	Droplet Precautions
Recent Exposure to Communicable Disease, Chicken Pox	Airborne and Contact Precautions
Recent Exposure to TB	Airborne Precautions
Active Pulmonary TB	Airborne Precautions