

VCH Acute Care Antibiotic Resistant Organism (ARO) Screening		
СРО	MRSA	VRE
All acute care inpatient admissions (excluding mental health)	All acute care inpatient admissions (excluding mental health)	VRE screening discontinued
Preadmission clinic when inpatient stay is expected within 14 days	Preadmission clinic when inpatient stay is expected within 14 days	
<ul> <li>Overnight stay or regular treatment visits to an outpatient clinic (e.g., dialysis) in an international hospital in the past 12 months</li> <li>As requested by IPAC</li> </ul>	<ul> <li>international hospital in past 12 months</li> <li>Hemodialysis or chemotherapy in past 12 months</li> <li>Shelter, homeless, illegal drug use, group home, or correction facility in past 12 months</li> <li>As requested by IPAC</li> </ul>	
No	High risk units *Universal admission screens include: direct admit or transfer from another unit/facility	
As requested by IPAC	ICU & BTHA – weekly on Monday	
<ul> <li>Rectal/ostomy (fecal stained)**</li> <li>If applicable:</li> <li>Sputum/tracheostomy aspirate if coughing</li> <li>Urine if catheterized</li> <li>Open wounds**</li> </ul>	<ul> <li>Nares</li> <li>Perineum or rectal/ostomy (fecal stained)!**</li> <li>If applicable:</li> <li>Open wounds**</li> <li>** if CPO also required, collect one swab per site and add all AROs indicated</li> </ul>	
Contact Precautions for confirmed and suspected CPO. Private room for confirmed CPO and high risk suspected CPO. Droplet and Contact Precautions if patient has productive cough or if ventilated in ICU	Contact Precautions  Droplet and Contact Precautions if MRSA found in sputum/tracheostomy aspirate and	Routine Practices in all locations for both VRE infection and colonization
	All acute care inpatient admissions (excluding mental health)  Preadmission clinic when inpatient stay is expected within 14 days  Overnight stay or regular treatment visits to an outpatient clinic (e.g., dialysis) in an international hospital in the past 12 months  As requested by IPAC  No  As requested by IPAC  Rectal/ostomy (fecal stained)**  If applicable: Sputum/tracheostomy aspirate if coughing Urine if catheterized Open wounds**  Contact Precautions for confirmed and suspected CPO. Private room for confirmed CPO and high risk suspected CPO. Droplet and Contact Precautions if patient	All acute care inpatient admissions (excluding mental health)  Preadmission clinic when inpatient stay is expected within 14 days  Overnight stay or regular treatment visits to an outpatient clinic (e.g., dialysis) in an international hospital in the past 12 months  As requested by IPAC  No  Rectal/ostomy (fecal stained)**  Rectal/ostomy (fecal stained)**  Sputum/tracheostomy aspirate if coughing  Urine if catheterized Open wounds**  Contact Precautions for confirmed and suspected CPO. Private room for confirmed CPO and high risk supected CPO. Droplet and Contact Precautions if patient

Refer to the <u>VCH Infection Control Admission Screening Tool</u>